ELSEVIER

Applied Nursing Research 19 (2006) 225–226

Applied Nursing Research

www.elsevier.com/locate/apnr

Author Index

Agrawal, S., 167
Alvine, C., 102
Anthony, M., 149

Badzek, L.A., 56
Baernholdt, M., 48
Barnason, S.A., 31
Beebe, L.H., 43
Berg, J., 182
Bertsch, C.M., 10
Blank, F.S., 70
Brey, B., 31
Briner, W., 167

Catlin, S., 31
Chaffin, M., 10
Chan, W., 16
Chang, C.-C., 221
Chang, S., 177
Chaudhury, H., 118
Chen, C.-Y., 191
Chung, C., 177
Clarke, S.P., 48
Corser, W.D., 63
Crogan, N.L., 102
Cunningham, R.S., 126

Davidson, P.M., 38
Davis, L.L., 212
De Bacquer, D., 156
Defloor, T., 156
Deiriggi, P., 56
Delicath, T., 149
Dick, M.J., 144
Doering, L., 182

Elswick Jr., R., 2
Emıroğlu, O.N., 134

Fiandt, K., 167
Fitzpatrick, J.J., 1, 55, 117, 173
Flowers, J., 144
Frazier, L., 16
Friend, L., 56
Fuentes, F., 16
Fung, N.I., 182

Gawlinski, A., 70, 182
Glazewki, L., 78
Glover, J., 56
Gray, D.P., 2
Grypdonck, M.H., 156

Habermann, B., 212
Halcomb, E.J., 38
Hannah, S.H., 144
Hanrahan, K., 95
Hayes, N., 171
Henneman, E.A., 70, 182
Henneman, P.L., 70
Hoyt, P., 110
Hughes, R.G., 223

Jang, H., 88

Kachorek, L., 78
Kearney, M.H., 163
Kıtış, Y., 134
Kleiber, C., 95
Kreman, R., 167
Kwak, S.A., 22

Lake, E.T., 51, 113
Lange, I., 220
Lee, J.B., 22
Leonhardy, K., 220
Leslie, N., 56
Lewallen, L.P., 144
Li, D.T., 216
Liehr, P., 16
Lim, J., 88
Lutgendorf, S., 95

Mackenzie, C.S., 105
Mahmood, A., 118
McCain, N.L., 2

0897-1897/$ – see front matter
doi:10.1016/S0897-1897(06)00123-6

McCarthy, A.M., 95
McClerking, C., 78
McCutcheon, H., 204
McDade, E., 2
Meininger, J.C., 16
Milisen, K., 171
Moore, S.M., 149
Mullins, S.M., 10

Narsavage, G., 149
Nieveen, J.L., 31

Park, S., 177
Patak, L., 182
Peterhans, B., 112
Poulin, P.A., 105
Powell, W., 144
Puntillo, K., 216

Representing the European Nursing Academy for Care of Older Persons (ENACO), 171
Robins, J.L.W., 2

Schuurmans, M.J., 171
Schwertfeger, R.U., 56
Scisney-Matlock, M., 78
Seidman-Carlson, R., 105
Shin, K.R., 22
Shin, Y., 88
Shurmur, S., 167
Simonelli, M.C., 163
Smalling, S., 16
Strumpf, N.E., 172

Thanavaro, J.L., 149
Tsai, Y.-F., 191
Tsalikian, E., 95

Valente, M., 118
Vanderwee, K., 156
Vogler, R., 16

Wall, Y., 144
Walter, J.M., 2
Wynd, C.A., 197

Yates, B.C., 167
Yi, H.R., 22
Yun, S., 88

Zeitz, K., 204
Zickefoose, K.T., 144
Zimmerman, L.M., 31

ELSEVIER

Applied Nursing Research 19 (2006) 227–230

Applied Nursing Research

www.elsevier.com/locate/apnr

Subject Index

Acupuncture, hand, and moxibustion, effectiveness in decreasing pain and "coldness" after hysterectomy in Korean women, 22
Adaptation, to diabetes, effects of home monitoring by public health nurse, 134
Adults
  Korean, with chronic diseases, program for exercise promotion among, 88
  with Parkinson's disease, feasibility of a psychoeducational intervention for caregivers of, 212
  smokers, relation of smoking patterns and beliefs to healthy behaviors among, 197
Advanced care planning, home health nurses' knowledge, comfort levels, barriers and personal participation in, 56
Advanced practice nurses, oncology, use of clinical practice guidelines, 126
AJN. *See American Journal of Nursing*
American Journal of Nursing, nurse editors' response to ANA's decision to discontinue affiliation with, 173
American Nurses Association, decision to discontinue affiliation with the *AJN*, response of nurse editors on, 173
ANA. *See* American Nurses Association
Auditory hallucinations, among schizophrenic Taiwanese patients, self-care symptom management for, 191

Behaviors, healthy, relation to smoking patterns and beliefs in adults, 197
Body fat percentage, in outpatients with schizophrenia, 43
Body mass index, in outpatients with schizophrenia, 43
BrAS. *See* Breast-Feeding Attitude Scale
Breast-Feeding Attitude Scale, to identify women at risk for early breast-feeding attrition, 144

Caregivers, of adults with Parkinson's disease, feasibility of a psychoeducational intervention for, 212
CHD. *See* Coronary heart disease
Children, strategies for salivary cortisol collection and analysis in, 95
Chronic diseases, program for exercise promotion among Korean adults with, 88
Clinical practice guidelines, use by oncology advanced practice nurses, 126
Coldness, and pain, after hysterectomy in Korean women, effectiveness of hand acupuncture and moxibustion for reduction of, 22
Commentary
  on need for evidence-based nursing practice in Latin America, 221
  on Problem Solving for Better Health Nursing program, 112
  on urgent leadership in nursing care for older persons, 172
Communication boards, in critical care, patients' perceptions on, 182
Computers, internet research in an international context, 48
Coronary heart disease, predictors of health promotion behavior in women without prior history of, 149
Cortisol, salivary, strategies for collection and analysis in children, 95
Critical care, patients' perceptions regarding communication boards in, 182

Data management, necessity of verbatim transcription of interview data, 38
Diabetes mellitus, effects of home monitoring by public health nurse on adaptation and self-care of patients with, 134
Diet, DASH, messages for hypertension treatment, development and evaluation of, 78
Drug exposure, prenatal. *See* Prenatal drug exposure

Editorials
  continued gap between research and professional practice, 1
  nurse editors' response to ANA's decision to discontinue affiliation with *AJN*, 173
  research
    on nursing languages, 117
    priorities, 223
  scientific misconduct among authors, 55
Elderly persons
  urgent need for leadership in nursing care for, 171
    commentary on, 172
  veterans, perceptions of their postdischarge outcome experiences, 63

0897-1897/$ – see front matter
doi:10.1016/S0897-1897(06)00135-2

Emergency department, academic, nurses' strategies to recover medical errors in, 70
Errors, medical, in an academic emergency department, nurses' recovery strategies of, 70
Erythema, nonblanchable, interrater reliability of trained nurses *vs.* researcher in using finger and transparent disk methods for identification of, 156
Evidence-based practice
  continued gap between research and professional practice, 1
  nursing, in Latin America, need for, 220
    commentary on, 221
Exercise, program for promotion among Korean adults with chronic diseases, 88

Finger method, *vs.* transparent disk method, reliability for NBE identification, 156

Growth patterns, and health service use among older siblings of infants with prenatal drug exposure, 10

Hallucinations, auditory, among schizophrenic Taiwanese patients, self-care symptom management for, 191
Hand acupuncture, and moxibustion, effectiveness in decreasing pain and "coldness" after hysterectomy in Korean women, 22
Health outcomes, multilevel models in research
  statistical and analytical issues, 113
  theory, design and measurement, 51
Health parameters, of outpatients with schizophrenia, description of, 43
Health promotion behavior, in women without prior history of CHD, predictors of, 149
Health services, use, and growth patterns among older siblings of infants with prenatal drug exposure, 10
Healthy behaviors, relation to smoking patterns and beliefs in adults, 197
HIV disease, tai chi as stress management for individuals with, 2
Home health nurses, in advanced care planning, knowledge, comfort levels, barriers and personal participation of, 56
Home monitoring, by public health nurse, effects on self-care and adaptation of diabetes patients, 134
Hospital discharge, older veterans' perceptions of their outcome experiences after, 63
Hyperlipidemia, effects of a motivational intervention on physiological outcomes in, 167
Hypertension
  development and evaluation of DASH diet tailored messages for treatment of, 78
  stage I, adding story-centered care to standard lifestyle intervention for people with, 16
Hysterectomy, decreased pain and "coldness" after, effectiveness of hand acupuncture and moxibustion in Korean women, 22

ICU. *See* Intensive care unit
Individual Nutrition Rx, assessment process, testing among nursing home residents, 102
Infants, with prenatal drug exposure, health services use and growth patterns among older siblings of, 10
INRx. *See* Individual Nutrition Rx
Intensive care unit, prevalence, intensity and relationships of coexisting symptoms in patients, 216
Internet, research in an international context, 48
Interview, necessity of verbatim transcription of, 38

Knowledge, of home health nurses in advanced care planning, 56
Koreans
  adult, with chronic diseases, program for exercise promotion among, 88
  women
    context of barriers to Pap testing in, 177
    effectiveness of hand acupuncture and moxibustion in decreasing pain and "coldness" after hysterectomy in, 22

Latin America, need for evidence-based nursing practice in, 220
  commentary on, 221
Leadership, in nursing care for older persons, urgent need for, 171
  commentary on, 172
Lifestyle intervention, standard, adding story-centered care to, for people with stage I hypertension, 16

Medical errors, in an academic emergency department, nurses' recovery strategies of, 70
Medical-surgical unit, nurses' perception advantages and disadvantages of single-occupancy *vs.* multioccupancy rooms in, 118
Mindfulness-based stress reduction intervention, for nurses and nurse aides, 105
Motivational intervention
  effects on physiological outcomes among hyperlipidemic persons, 167
  nurse-delivered, feasibility and efficacy in promoting postpartum weight loss, 163
Moxibustion, and hand acupuncture, effectiveness in decreasing pain and "coldness" after hysterectomy in Korean women, 22
Multioccupancy room, *vs.* single-occupancy room, in medical-surgical units, nurses' perception of advantages and disadvantages, 118

NBE. *See* Nonblanchable erythema
Nonblanchable erythema, interrater reliability of trained nurses *vs.* researcher in using finger and transparent disk methods for identification of, 156
Nurse aides, mindfulness-based stress reduction intervention for, 105

Nurse editors, response to ANA's decision to discontinue affiliation with *AJN*, 173
Nurses
  home health, in advanced care planning, knowledge, comfort levels, barriers and personal participation of, 56
  mindfulness-based stress reduction intervention for, 105
  motivational intervention delivered by, feasibility and efficacy in promoting postpartum weight loss, 163
  perceptions of advantages and disadvantages of single-occupancy room *vs.* multioccupancy room in medical-surgical units, 118
  trained, *vs.* researcher, interrater reliability in observing NBE using finger and transparent disk methods, 156
Nursing, evidence-based, in Latin America, need for, 220
  commentary on, 221
Nursing care, for older persons, urgent need for leadership in, 171
  commentary on, 172
Nursing homes, residents, testing of INRx assessment process among, 102
Nursing languages, research on, 117

Older persons. *See* Elderly persons
Oncology advanced practice nurses, use of clinical practice guidelines, 126
Outcomes
  health, multilevel models in research
    statistical and analytical issues, 113
    theory, design and measurement, 51
  physiological, among hyperlipidemic persons, effects of motivational intervention on, 167
  postdischarge, veterans' perceptions of, 63
Outpatients, with schizophrenia, description of health parameters of, 43

Pain, and "coldness," after hysterectomy in Korean women, effectiveness of hand acupuncture and moxibustion, 22
Pap testing, in Korean women, context of barriers to, 177
Parkinson's disease, feasibility of a psychoeducational intervention for caregivers of adults with, 212
Participation, of home health nurses in advanced care planning, 56
Patient perception, regarding communication boards in critical care, 182
Patient surveillance, impact of frequent collection of postoperative vital signs on detection of complications, 204
PCI. *See* Percutaneous coronary intervention
Percutaneous coronary intervention, recovery patterns following, 31
Physical health, parameters, of outpatients with schizophrenia, description of, 43
Postoperative period, impact of frequent collection of vital signs on detection of complications, 204
Prenatal drug exposure, infants with, health services use and growth patterns among older siblings of, 10
Prevalence, and intensity and relationships, of coexisting symptoms in ICU patients, 216
Problem Solving for Better Health Nursing, 110
  commentary on, 112
Professional practice, and research, continued gap between, 1
Psychiatric health, parameters, of outpatients with schizophrenia, description of, 43
Psychoeducational intervention, for caregivers of adults with Parkinson's disease, feasibility of, 212
Psychoneuroimmunology, tai chi as stress management for individuals with HIV disease, 2
Public health nurse, home monitoring by, effects on self-care and adaptation of diabetes patients, 134

Recovery patterns, following PCI, 31
Research
  adding story-centered care to standard lifestyle intervention for people with stage I hypertension, 16
  assessment of home health nurses in advanced care planning, 56
  BrAS to identify women at risk for early breast-feeding attrition, 144
  clinical practice guidelines use by oncology advanced practice nurses, 126
  context of barriers to Pap testing in Korean women, 177
  describing health parameters of outpatients with schizophrenia, 43
  development and evaluation of DASH diet tailored messages for hypertension treatment, 78
  effectiveness of hand acupuncture and moxibustion in decreasing pain and "coldness" in Korean women who had hysterectomy, 22
  effects of home monitoring by public health nurse on self-care and adaptation of diabetes patients, 134
  effects of motivational intervention on physiological outcomes among hyperlipidemic persons, 167
  feasibility and efficacy of a nurse-delivered motivational intervention in promoting postpartum weight loss, 163
  feasibility of a psychoeducational intervention for caregivers of adults with Parkinson's disease, 212
  health outcomes, multilevel models in
    statistical and analytical issues, 113
    theory, design and measurement, 51
  health services use and growth patterns among older siblings of infants with prenatal drug exposure, 10
  impact of frequent collection of postoperative vital signs on detection of complications, 204
  internet-based, in an international context, 48
  interrater reliability between researcher and trained nurses in observing NBE using finger and transparent disk methods, 156

Research (continued)
mindfulness-based stress reduction intervention for nurses and nurse aides, 105
necessity of verbatim transcription of interview data, 38
need for evidence-based nursing practice in Latin America, 220
commentary on, 221
nurse editors' response to ANA's decision to discontinue affiliation with *AJN*, 173
nurses' perception of advantages and disadvantages of single-occupancy *vs.* multioccupancy rooms in medical-surgical units, 118
nurses' strategies to recover of medical errors in an academic emergency department, 70
nursing languages, 117
older veterans' perceptions of their postdischarge outcome experiences, 63
patients' perception regarding communication boards in critical care, 182
predictors of health promotion behavior in women without prior history of CHD, 149
prevalence, intensity and relationships of coexisting symptoms in ICU patients, 216
priorities in, 223
Problem Solving for Better Health Nursing program, 110
commentary on, 112
and professional practice, continued gap between, 1
program for exercise promotion among Korean adults with chronic diseases, 88
on psychoneuroimmunology, tai chi as stress management for individuals with HIV disease, 2
recovery patterns following PCI, 31
relation of smoking patterns and beliefs to healthy behaviors among adult smokers, 197
scientific misconduct among authors, 55
self-care symptom management for auditory hallucinations among schizophrenic Taiwanese patients, 191
strategies for salivary cortisol collection and analysis in children, 95
testing of INRx assessment process among nursing home residents, 102
urgent need for leadership in nursing care for older persons, 171
Researcher, *vs.* trained nurses, interrater reliability in observing NBE using finger and transparent disk methods, 156
Risk, for early breast-feeding attrition in women, BrAS for identification of, 144

Salivary cortisol, strategies for collection and analysis in children, 95
Schizophrenia
describing health parameters of outpatients with, 43
self-care symptom management for auditory hallucinations among Taiwanese patients with, 191
Self-care
of diabetes patients, effects of home monitoring by public health nurse, 134
symptom management, for auditory hallucinations, among schizophrenic Taiwanese patients, 191
Siblings, older, of infants with prenatal drug exposure, health services use and growth patterns among, 10
Single-occupancy room, *vs.* multioccupancy room, in medical-surgical units, nurses' perception of advantages and disadvantages, 118
6-Minute walk test, in outpatients with schizophrenia, 43
Smoking, patterns and beliefs, relation to healthy behaviors among adult smokers, 197
Story-centered care, adding to standard lifestyle intervention, for people with stage I hypertension, 16
Stress management, tai chi for individuals with HIV disease, 2
Stress reduction, mindfulness-based intervention for nurses and nurse aides, 105
Symptom management, self-care, for auditory hallucinations, among schizophrenic Taiwanese patients, 191
Symptoms, coexisting, in ICU patients, prevalence, intensity and relationships of, 216

Tai chi, as stress management for individuals with HIV disease, 2
Taiwanese, schizophrenic patients, self-care symptom management for auditory hallucinations among, 191
Transcription, verbatim, of interview data, necessity of, 38
Transparent disk method, *vs.* finger method, reliability for NBE identification, 156

Verbatim transcription, of interview data, necessity of, 38
Veterans, older, perceptions of their postdischarge outcome experiences, 63
Vital signs, postoperative, impact of frequent collection on detection of complications, 204

Weight loss, postpartum, feasibility and efficacy of a nurse-delivered motivational intervention in promotion of, 163
Women
Korean, context of barriers to Pap testing in, 177
without prior history of CHD, predictors of health promotion behavior in, 149
at risk for early breast-feeding attrition, identified by BrAS, 144